

Task: GLO

Global warming

CEOI 2018, day 1. Available memory: 256 MB.

14.08.2018

Global warming is an important issue and Johnny knows about it. He decided to make an analysis of historical temperatures and find a subsequence of days (not necessarily consecutive) where the temperature was strictly increasing. It will convince the non-believers!

Johnny has found historical data from n consecutive days. The temperature on the i -th day was t_i .

Formally, we are interested in finding the length of the longest increasing subsequence (LIS) of (t_1, t_2, \dots, t_n) , that is, the largest possible k for which it is possible to choose an increasing sequence of indices $1 \leq a_1 < a_2 < \dots < a_k \leq n$ such that $t_{a_1} < t_{a_2} < \dots < t_{a_k}$.

Johnny wants to find a really long subsequence and that is why he decided to cheat a bit. He will first choose a non-empty interval of days and an integer d ($-x \leq d \leq x$) and he will increase the temperature in each of those days by d . A small change like that probably will not be noticed by the community, while at the same time it should make the LIS longer. It is allowed to choose $d = 0$.

What is the largest possible length of the LIS after a change?

Input

The first line of the standard input contains two space-separated integers n and x ($1 \leq n \leq 200\,000$, $0 \leq x \leq 10^9$), the number of days and the limit for the absolute value of d .

The second line contains n integers t_1, t_2, \dots, t_n ($1 \leq t_i \leq 10^9$) separated by spaces, the sequence of historical temperatures.

Output

Print one integer, the largest possible length of the LIS after a single change.

Example

For the input data:

8 10
7 3 5 12 2 7 3 4

the correct result is:

5

Explanation of the example: Johnny can choose an interval $[2, 3]$ and $d = -5$, which means decreasing t_2 and t_3 by 5. In this case the new sequence is $(7, -2, 0, 12, 2, 7, 3, 4)$, where he can find a LIS $(-2, 0, 2, 3, 4)$. The length of the LIS is 5.

Grading

The test set is divided into the following subtasks with additional constraints. Tests in each of the subtasks consist of one or more separate test groups. Each test group may contain one or more test cases.

Subtask	Constraints	Points
1	$n, x \leq 10$	5
2	$n, x \leq 50$	10
3	$n \leq 1000$	13
4	$x = 0$	10
5	$x \leq 5, n \leq 50\,000$	20
6	$x = 10^9$	17
7	no additional constraints	25